


Drugs and the Brain

A Beginner's Guide to Stimulants and Empathogens



Version: 1.3

Original version: 18/03/2014

Revision date: 03/04/2014

Part 1: The lizard brain^{1,2}

The brain controls almost everything we do: from our first breath in the morning to our dreams at night. Very little that we do, think or feel takes place without the brain being in charge.

Much of what the brain does happens on an automatic or *subconscious* level. For example, we blink many times a day without consciously thinking about it, but it is our brain that makes us do this (to clean, protect and moisten our eyeballs).

Buried deep within the human subconscious brain (also called the lizard brain) are survival instincts:

- To eat (to stay alive)
- To reproduce (to continue the species)
- To respond to threats or aggression (to protect yourself, your family and what you own)





Image credits: see references section at the end of this document

Drugs and the Brain: A Beginner's Guide

There are certain areas in our brain whose job it is to make us feel good when we do these things, so when we eat, have sex or take exercise our body releases chemicals that make us feel happy or excited. These parts of the brain are connected in networks called **reward pathways** because they release **reward chemicals** when we do something that makes us more likely to survive.

The picture to the right shows two of the brain's reward pathways. There are many of these pathways in the brain; they are like phone or computer networks with information flying along wires.

Drugs that get you high use these reward pathways to make us feel good. They are like pirates taking over a ship: they hijack these parts of the brain and make them release lots of reward chemicals, which is why we feel high or stoned.




However, if we keep taking these drugs again and again they can take over the reward pathways in the brain until all we want to do is to take the drug. Nothing else makes us feel good any more apart from getting high and this can become a habit.

Once you get to this stage:

- You might have cravings
- You start to want to take the drug more than doing anything else
- You find yourself keeping using even when you want to stop
- You become more moody and irritable when you can't get hold of it.

Part 2: The reward pathways ^{2, 3, 4, 6, 7}

The brain is a big network made partly of **neurons** (nerve cells). It uses chemicals called **neurochemicals** (nerve chemicals) to send and receive signals between these neurons.




In this image you can see detail of how signals pass between neurons. This is how the brain communicates: every time we have a thought or feeling this is because of neurochemicals passing between neurons.

Drugs and the Brain: A Beginner's Guide

The picture to the right is a simple version of part of the brain's reward pathway that controls some of our moods and attitudes.

When this reward pathway is activated (for example by food, water or sex), information travels from the VTA to the Nucleus Accumbens and then up to the Prefrontal Cortex.



The VTA uses an important chemical called **dopamine**

to send this information. Dopamine controls things like what is important to us, what gives us pleasure, how we feel, how we learn and what motivates us. Dopamine gets stimulated by many drugs and is part of what makes you feel high, and how much you feel you want or need a drug. It is often linked to stimulants, so amphetamine (speed) and cocaine are typical drugs that raise dopamine levels. As the levels of dopamine increase in the brain, other areas get stimulated, including the Prefrontal Cortex. The Prefrontal Cortex is very important, especially where drugs are concerned, as it controls a number of different functions, including:

- Behaviour (*controlling your actions, acting in an appropriate way*)
- Development of conscience (*knowing right from wrong and doing the right thing*)
- Decision making (*gathering information before making a decision*)
- Impulse control (*being able to control your feelings*)
- Focus (*being able to stay on track with a project until it's finished*)

Drugs and the Brain: A Beginner's Guide

People who have problems with the Prefrontal Cortex can experience:

- Short attention span
- Problems with impulse control (*being impatient, 'kicking off'*)
- Poor organisation and planning skills
- Hyperactivity
- Emotional problems (*such as shutting down, withdrawing from people*)
- Misreading situations
- Being dishonest
- Problems with learning
- Short term memory problems
- Anxiety
- Serious mental health problems (*such as psychosis*)

Using drugs changes the way that this part of the brain grows, develops and works. Young people can develop problems with drugs more easily as the Prefrontal Cortex can still be growing and developing when people reach their early twenties.

Drugs and the Brain: A Beginner's Guide

Part 3: Neurochemicals ^{1, 2, 3, 4}

Dopamine is a neurochemical that the brain uses to send messages to its different parts. Different neurochemicals control different parts of the brain.




Neurochemicals are like the various types of messages found in a phone network, for example phone calls, texts, emails or instant messages.

Here are some more examples of neurochemicals:

Neurochemical	What it does
Noradrenaline	How awake or aroused you feel, attention span and confidence.
Adrenaline	Similar to noradrenaline and also controls energy.
Dopamine	Motivation, wants, pleasure, love and the reward pathway. This is a big part of what makes you want to take more of a drug.
Serotonin	Memory, emotions, wakefulness, sleep, mood (depression), sexual activity and temperature. It is associated with euphoria (happy feelings).
GABA	Controls impulses, muscle relaxation and arousal and slows down the brain. Plays a big part in sleep and 'feeling good'.
Endorphin	The senses, especially pain, emotional and physical distress. A lot of the high you get from exercise comes from endorphins.
Histamine	Involved in sleep and waking patterns and inflammatory response (for example allergic reactions).

Part 4: How a drug affects your mood ^{1, 2, 7, 8, 9, 10, 15}

When you do something that triggers the brain's reward pathway (for example, taking exercise), neurochemicals are released by the **transmitting neuron**.




Some of the neurochemicals land on the **receptors** on the receiving neuron, and this is what sends out feelings of pleasure to other parts of the brain. Most of the extra neurochemicals are absorbed back by travelling through the **reuptake transporter**, ready to be used again if necessary. Some neurochemicals do not get absorbed, and this is thought to play a part in **drug toxicity** (poisoning, or overdose).

We will now look at what happens when you take certain stimulant and empathogenic drugs.

Drugs and the Brain: A Beginner's Guide

Cocaine


When you take cocaine it affects the neurons in two ways. Firstly it stimulates the transmitting neuron to send out lots of dopamine. It also blocks the dopamine reuptake transporter which means that the extra dopamine that has been released ends up flying around your brain, triggering and stimulating the dopamine receptors which is why you feel high.



If you keep on using stimulant drugs over and over again you can use up all your dopamine (this is called dopamine depletion). This can lead to problems including depression and other mental health issues, issues with learning, hyperactivity, cravings, dependency or addiction and problems releasing dopamine in the future.

Amphetamine (speed)

Like cocaine, amphetamine increases the amount of dopamine in the neurons, but it works in a different way. The amphetamine molecule is similar in shape to dopamine and it can enter the transmitting neuron through its reuptake transporters. It then forces the dopamine molecules out through the transporters. This can be toxic, and can lead to life-threatening hyperthermia (overheating).



If you find yourself doing stimulant drugs too often you can become tired, run down, low in mood and possibly have mood swings and some paranoid thoughts. The best thing that you can do is to **take a break** and allow your body to recharge. Plenty of sleep and food can help your body and mind to recover.

Drugs and the Brain: A Beginner's Guide


MDMA (ecstasy)

Although MDMA causes the release of dopamine, another neurochemical it stimulates is called **serotonin**. Serotonin is an important neurochemical that helps you to feel relaxed and controls mood, appetite and sleep. It is serotonin that gives MDMA a lot of its 'loved up' feeling and this is why ecstasy is called an **empathogen** (empathy is when you place yourself in someone else's shoes and feel what they are feeling).

Dopamine pathways

Functions

- Reward (motivating)
- Pleasure, euphoria
- Motor function (fine-tuning)
- Compulsion
- Re-dosing



Serotonin pathways

Functions


- Mood
- Memory processing
- Sleep
- Cognition/thinking
- Appetite

When you take empathogens they release serotonin in your brain, but can leave your serotonin levels very low afterwards and can also lead to life-threatening hyperthermia. If you take more drugs to try to get that good feeling back you might find your body can't produce the high, a bit like trying to flush a toilet twice in a row!

Using up your stores of serotonin is partly the reason why you might feel down, angry, sad or upset a few days after taking empathogens. And it is not just low levels of serotonin that causes problems; very high levels can lead to a medical condition called Serotonin Syndrome which can be fatal (see page 13).

Drugs and the Brain: A Beginner's Guide

When you take MDMA it affects the neurons in two ways. Firstly it makes the transmitting neuron send out lots of serotonin. It also blocks the serotonin reuptake transporter, and it is this extra serotonin flooding the area between the neurons that leads to the 'loved up' feeling.




Drugs and the Brain: A Beginner's Guide

Mephedrone (bubble, mkat)

Mephedrone is a powerful drug that greatly increases both dopamine and serotonin levels, and in animal studies caused about a 500% increase in dopamine and a 900% increase in serotonin. This makes it both a stimulant drug and an empathogen. As the effects of the drug start to wear off, serotonin levels drop quite quickly which is why people want to take more and more of the drug (this is called 'redosing').

As mephedrone is a relatively new drug we are not completely sure what it does in the brain. It acts mostly like amphetamine, but also affects serotonin levels.



Serotonin syndrome¹⁶

This is a medical condition caused when your body releases too much serotonin. It can occur in a lot of different situations, one of these is when you take recreational drugs that effect your serotonin levels. It can kill you if not recognised and dealt with both quickly and effectively.

Symptoms include: agitation, anxiety, overheating, shivering, restlessness, twitching, high blood pressure, flushed skin, stomach noises, vomiting, diarrhea, dilated pupils and mental distress.

If someone you are with is showing some of these signs **call an ambulance immediately**.

While waiting for the ambulance do what you can to **cool the person down and calm them down**. It is dangerous to restrain people in this state.

For further information on Overdose & Emergencies see DrugWatch Information Sheet.

Drugs and the Brain: A Beginner's Guide

Part 5: How drug use can lead to dependency ^{2, 3, 4, 5, 7}

The drugs that have the strongest effect on your brain are stimulant drugs (like speed, cocaine and mephedrone) and opiate drugs (like heroin). The way you take a drug also affects how likely you are to become dependent. In general, the quicker and stronger the drugs affects you, the more your dopamine levels 'spike'. This is why injecting and smoking drugs can be more likely to lead to dependency than eating or swallowing them.

When you first start taking a drug you only need a relatively small amount to get you high. But if you keep on taking it your brain can get used to all the extra reward chemicals and you need to take more and more to feel that same high. This is called **tolerance**.

Your brain gets used to these extreme levels of reward chemicals, and so normal activities that you used to enjoy (like exercise, eating a nice meal or spending time with your family) no longer make you feel happy.

You might find yourself feeling **bored** the whole time... unless you are taking drugs. This is because your brain is getting used to the extreme 'spikes' in reward chemicals.

When you aren't taking drugs the levels of these chemicals drop and your brain starts wanting more.

It sends out warning messages around the brain and these messages get sent out to different parts of the body. You can feel these in different ways, some people feel edgy, angry, anxious, stressed, tearful, shaky, sweaty or as if they have a knot in their stomach... these feelings are called **cravings**.



Part 6: Cravings ^{11, 12}

Cravings can be very sneaky. They are the brain's way of trying to get you to take drugs so that it can balance out your reward chemicals. They often make you feel agitated, angry or stressed and it is very easy to believe that you are feeling upset with people when in fact it is just your brain wanting to get its happy chemicals back.

If you don't learn to **recognise your cravings** you can often behave in ways that might be out of character, for example getting into arguments with people and 'kicking off' for no reason. There are lots of reasons that people have cravings, and different people feel them in different ways. This is something that your drugs worker can help with, as we all react differently to cravings and have our own ways of coping with them.





Cravings can come in many forms: for example it can be as simple as thinking about drugs on Friday morning before you are due to go out that night, a very strong, tense, angry feeling if you can't get hold of any drugs when you were expecting them, the ache for more when your supplies are running low, or the tingle you might get in your stomach when you've been off drugs for a few days and think about them suddenly. These feelings are all caused by neurochemicals (mostly dopamine) being released in your brain.

Lots of things can make you crave drugs. These are called **triggers**. Triggers can be thoughts, feelings, sights, sounds, smells, places or situations. Examples could be: it's a hot day and you see a beer glass, or someone walks past you and you smell cannabis smoke.

Drugs and the Brain: A Beginner's Guide

When you use drugs regularly you train your brain to get a big reward when you take them. Think about how you feel when you are just about to buy drugs, and when you have just bought them. Your brain releases reward chemicals like dopamine even before you've taken the drug because you have trained it to know that it will soon get a big reward (see picture on the right).



When you have a craving your brain thinks it's going to get drugs, so it releases some reward chemicals. When it doesn't get the drugs it was expecting your levels of reward chemicals drop, and you can feel bad, stressed, frustrated, sick or angry.

If you don't learn to interrupt or control your cravings you can find them taking over your thoughts, and if you keep giving in to them it can become more and more difficult to resist taking drugs.


Part 7: What is dependency? ^{5, 6, 13, 14}

Being dependent on a drug is when you feel you *have* to take it, when you can't stop or cut down, and when you keep on using even when you know the drug is causing you problems.

If we think back to the reward system in the brain. Firstly you are motivated to do something for survival (for example, eat) and then your brain releases chemicals which are your reward. This ends the reward process.

Motivation → **Reward Chemicals** → **Reward**
food, career, sex etc.

When we take drugs we change the way this pathway works; taking drugs give us 'free' reward chemicals. This can cause problems as it can create a loop which can lead to dependency, when the drugs become the motivation and survival instinct.



Dependency is when drugs take control over your behaviour. It usually begins with casual or recreational use ('every now and then' or 'just at weekends') and ends when you feel you have to take them.

Some people can use drugs without problems, and others become dependent very easily. There are many reasons why this might happen: it might be something in your brain, something in your personality, something in your upbringing or maybe a combination of these things. At the moment, no one knows why.

Part 8: Are drugs becoming a problem? ¹⁷

Working this out isn't simple, and admitting it isn't easy because we all want to think we're in control. Below are some of the things that can suggest that your drug use is becoming a problem. If you have answered yes to three or more of the following questions in the past year it suggests your drug use might be becoming a problem and that it might help to speak to someone about it.

Do you need more to get high, or get the effect you wanted?

Do you have any signs of physical withdrawal, or do you find yourself thinking about a lot when you don't have it?

Do you find yourself taking more in a session than you'd originally planned, or find yourself taking it for longer sessions?

Do you find yourself wanting to cut down or stop taking ?

Do you spend a lot of time scoring or recovering from a session?

Have you stopped, or cut down activities such as school, college, training, work, hanging out with friends or family because of your use of?

Have you kept on using despite knowing that it is causing ongoing problems to your physical or mental health?

Has using caused you troubles with the law (police, YOT, probation)?

References

- ¹ *Drug Addiction and Its Underlying Neurobiological Basis: Neuroimaging Evidence for the Involvement of the Frontal Cortex*. Goldstein, R. & Volkow, N. *Am J Psychiatry*. 2002 October; 159(10): 1642–1652. Accessed online 24/01/2014 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1201373>
- ² *Neural systems of reinforcement for drug addiction: from actions to habits to compulsion*. Everitt B. & Robbins, T. *Nature Neuroscience* 8, 1481 - 1489 (2005). Published online: 26 October 2005 | doi:[10.1038/nn1579](https://doi.org/10.1038/nn1579). Accessed online 24/01/2014: <http://www.nature.com/neuro/journal/v8/n11/full/nn1579.html>
- ³ *Addiction, compulsive drug seeking, and the role of frontostriatal mechanisms in regulating inhibitory control*. Feil, J., Sheppard, D., Fitzgerald, P., Yücel, M., Lubman, D. & Bradshaw, J. *Neuroscience & Biobehavioral Reviews* Volume 35, Issue 2, November 2010, Pages 248–275. Accessed online 24/01/2014: <http://www.sciencedirect.com/science/article/pii/S0149763410000461>
- ⁴ *Pleasure systems in the brain*. Bozarth, M. (1994). In D.M. Warburton (ed.), *Pleasure: The politics and the reality* (pp. 5-14 + refs). New York: John Wiley & Sons. Accessed online 24/01/2014: <http://wings.buffalo.edu/aru/ARUreport01.htm>
- ⁵ *The neurobiology of pleasure, reward processes, addiction and their health implications*. Esch, T. & Stefano, G. *Neuroendocrinology Letters* No.4 August Vol.25, 2004. Copyright © 2004 Neuroendocrinology Letters ISSN 0172–780X www.nel.edu. Accessed online 24/01/2014: http://www.nel.edu/pdf/_NEL250404R01_Esch-Stefano_p_pdf
- ⁶ *Dynamic mapping of human cortical development during childhood through early adulthood*. Gogtay, N., Giedd, J., Lusk, L., Hayashi, K., Greenstein, D., Vaituzis, A., Nugent, T., Herman, D., Clasen, L., Toga, A., Rapoport, J. & Thompson, P. *Proc Natl Acad Sci USA*. 2004 May 25; 101(21): 8174–8179. Published online 2004 May 17. Accessed online 24/01/2014: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC419576/>
- ⁷ *Brain development during childhood and adolescence: a longitudinal MRI study*. Giedd, J., Blumenthal, J., Jeffries, N., Castellanos, F., Liu, H., Paus, T., Evans, A. & Rapoport, J. *Nature Neuroscience* Volume 2 No 10, October 1999 © 1999 Nature America Inc. • <http://neurosci.nature.com>. Accessed online 24/01/2014: http://www.math.tau.ac.il/~dms/Longitudinal/brain_MRI.pdf
- ⁸ *The Brain From Top To Bottom*. Online Resource <http://thebrain.mcgill.ca/> Accessed 24/01/2014.
- ⁹ *Mephedrone, compared with MDMA (ecstasy) and amphetamine, rapidly increases both dopamine and 5-HT levels in nucleus accumbens of awake rats*. Kehr, J., Ichinose, F., Yoshitake, S., Gojny, M., Sievertsson, T., Nyberg, F. & Yoshitake, T. *Br J Pharmacol*. 2011 December; 164(8). Accessed online 24/01/2014: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1201373/>
- ¹⁰ *Designer cathinones — An emerging class of novel recreational drugs*. Zawilska, J. & Wojcieszak, J. *Forensic Science International* Volume 231, Issue 1, Pages 42-53, 10 September 2013. Accessed online 24/01/2014: <http://download.journals.elsevierhealth.com/pdfs/journals/0379-0738/PIIS0379073813002387.pdf>
- ¹¹ *Alcohol Craving and Relapse Prediction*. Heinz, A., Beck, A., Mir, J., Grüsser, S., Grace, A., Wrase, J. *Advances in the Neuroscience of Addiction*. 2nd edition. Kuhn CM, Koob GF, editors. Boca Raton (FL): CRC Press; 2010. Accessed online 24/01/2014: <http://www.ncbi.nlm.nih.gov/books/NBK53355/>
- ¹² *A neural substrate of prediction and reward*. Schultz, W., Dayan P. & Montague, P. (1997) *Mar* 14; 275(5306):1593-9. Accessed online 24/01/2014 <http://www.gatsby.ucl.ac.uk/~dayan/papers/sdm97.pdf>
- ¹³ *Neurobiology of Alcohol Dependence: Focus on Motivational Mechanisms*. Gilpin, N. & Koob, G. *National Institutes of Health Online Publications*. Accessed 24/01/2014: <http://pubs.niaaa.nih.gov/publications/ahr313/185-195.htm>
- ¹⁴ *Addiction Is a Brain Disease, and It Matters*. Leshner, A. *Science* 278, 45 (1997); DOI: [10.1126/science.278.5335.45](https://doi.org/10.1126/science.278.5335.45). Accessed 24/01/2014: <http://www.ecnp.eu/~media/Files/ecnp/communication/talk-of-the-month/Wim%20van%20den%20Brink/Addiction%20is%20a%20brain%20disease%20and%20it%20matters.pdf>
- ¹⁵ *A Marked inhibition of mesolimbic dopamine release: a common feature of ethanol, morphine, cocaine and amphetamine abstinence in rats*. Rossetti, Z., Hmaidan, Y., Gessa, G. & Brodie, B. Department of Neuroscience, University of Cagliari, Italy. *European Journal of Pharmacology* (Impact Factor: 2.59). 11/1992; 221(2-3):227-34. DOI:10.1016/0014-2999(92)90706-A Source: PubMed.
- ¹⁶ *Serotonin Syndrome*. Volpi-Abadie, J., Kaye, A. & Kaye, A. (2013). *The Ochsner Journal* 13:533-540. Academic Division of Ochsner Clinic Foundation.
- ¹⁷ Definitions of substance dependence and substance abuse adapted from the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (Copyright 2000, American Psychiatric Association).

Image credits

- Page 1 *Lizard*: Original image Gallotia Simonyi, Female, Hierro giant lizard. Scientific illustration, Pen (2003). <http://www.norarte.es> Permission is granted to copy, distribute and/or modify this document under the terms of the [GNU Free Documentation License](http://www.gnu.org/licenses/old-licenses/gpl-2.0.html). Accessed online 24/01/2014 http://commons.wikimedia.org/wiki/File:Gallotia_simonyi-female_norarte.jpg. Revised in Adobe Illustrator and Photoshop by Mark Adley.
- Pages 2, 4 & 10 *Brain pathways*: Mark Adley, based on dopamine and serotonin pathways image. Original source: <http://www.drugabuse.gov/pubs/teaching/largegifs/slide-2.gif> (2009), NIH [Public Domain] Accessed online 24/01/2014 http://commons.wikimedia.org/wiki/File:Dopamine_and_serotonin_pathways.gif
- Page 2 *Pirate*: Pyle, Howard; Johnson, Merle De Vore (ed) (1921) *With the Buccaneers in Howard Pyle's Book of Pirates: Fiction, Fact & Fancy Concerning the Buccaneers & Marooners of the Spanish Main*, New York, United States, and London, United Kingdom: Harper and Brothers, pp. Plate facing p. 76 Retrieved on 14 April 2010. Howard Pyle [Public Domain] Accessed online 24/01/2014 http://commons.wikimedia.org/wiki/File:Pyle_pirates_burying2.jpg
- Page 3 *Neuron*: Purchased clip art, revised in Adobe Illustrator by Mark Adley.
- Pages 3, 7, 8, 9, 11 & 12 *Synapses*: Images created in Adobe Illustrator by Mark Adley. Based on images from *The Brain Top To Bottom* <http://thebrain.mcgill.ca/>. Accessed 24/01/2014. The content of the site *The Brain from Top to Bottom* is under copyleft. The concept of “copyleft” is a method of providing free access to the results of original work and of encouraging people to reproduce and even modify this work on an equally free basis. http://thebrain.mcgill.ca/flash/pop/pop_copy/pop_copy_i.html
- Page 6 *Network*: converted to vector and edited in Adobe Illustrator by Mark Adley. Based on the image <http://odak.co.uk/business-users/virtual-private-networks-vpn>. Kind permission to use the image was given by email, subject RE: Message from ODAK webpage form, sent by Aytekin Yelden [aytekin@odak-online.com] Tue 21/01/2014 17:01.
- Page 13 *Bored*: converted to vector and edited in Adobe Illustrator by Mark Adley. Based on the image <http://commons.wikimedia.org/wiki/File:Bored.png>. Permission is granted to copy, distribute and/or modify this document under the terms of the [GNU Free Documentation License](http://www.gnu.org/licenses/old-licenses/gpl-2.0.html). Accessed online 24/01/2014.
- Page 14 *Cravings*: Masked man (2006) by Liftarn. Source <http://openclipart.org/c/ghost/media/files/liftarn/1243>. This file is from the Open Clip Art Library, which released it explicitly into the public domain. Accessed online 24/01/2014 http://commons.wikimedia.org/wiki/File:Masked_man.svg

Drugs and the Brain: A Beginner's Guide

This guide was written and designed by Mark Adley, in collaboration with UK DrugWatch and N2L. Mark is a drugs worker in the North East of England and the creator of the **Drugs Wheel**, a new psychoactive substance awareness model (available at www.thedrugswheel.com) designed to illustrate the wide range of new substances that are available.



N2L (Never Too Late) is a confidential service for young people under 18 who live in North Tyneside and are experiencing difficulties in relation to the use of drugs and alcohol.

Phone: (0191) 643 8802, Fax: (0191) 643 8801

Email: n2l@northtyneside.gov.uk



UK DrugWatch is an informal online professional information network established by a group of professionals working in the UK drugs sector. The aim of the group is to raise/establish standards for drug information, alerts and warnings. It is currently an unfunded, bottom-up initiative that works in the spirit of mutual co-operation. A list of current members, and a selection of drug briefings can be found [here](#).



A special thanks to Colin Davidson, neurophysiologist from TicTac Communications, Ltd., for invaluable advice.

Drug and Alcohol Monitoring and Information System (DAMIS)

Activity report

October 2014 - March 2015

Background

DAMIS is an “early warning system” designed to find out about emerging trends in drug and alcohol misuse, so we can act quickly and provide relevant information or advice to those who misuse drugs or alcohol.

The kind of information DAMIS collects includes:

1. A sudden increase in a particular drug being misused
2. Drugs being misused in new ways
3. New drugs becoming available
4. Emergence of substances with unexpected unpleasant or dangerous effects

The DHSSPS oversee DAMIS with support from the lead partners: the Public Health Agency, the Department of Justice and the Police Service of Northern Ireland.

As at April 2015, DAMIS has an email circulation list of 285 individuals. Unless otherwise stated, information sent out by DAMIS is emailed to all those on the list. People can be added to this list by emailing a request to damis@hscni.net.

Anyone can send in information to DAMIS. It should be emailed to damis@hscni.net.

The following tables summarize key information received by DAMIS, and circulated by DAMIS, in the 6 months from October 2014 to March 2015. If you would like further information on anything listed here, please contact damis@hscni.net

Information received by DAMIS during October 2014 – March 2015

Information received by DAMIS is shown in the table below:

Emails received by DAMIS October 2014 – March 2015	
Date	Topic
20 th October	"Trippy" synthetic cannabinoid widely used in Southern Trust area, with psychological withdrawal extreme and longer-lasting than other synthetic drugs.
5 th November	Addiction Services client with polydrug use (including injecting mephedrone and diphenidrine) had a stroke.
11 th November	Drug sold as a synthetic cannabinoid, believed to be called "sky blue", linked to a number of people staying in a hostel in Belfast requiring emergency medical treatment
12 th November	Relates to above. Drug may be called "sky high" or "sky's the limit"
24 th November	Users report pills sold as Ecstasy on the streets in Derry, with the names <i>Green Heineken</i> , <i>Gold Bar</i> , and <i>Chubba Chubba</i> have led to higher levels of anxiety / depression than expected (more information was requested)
28 th November	White heroin sold as cocaine in Amsterdam
10 th December	Unknown stimulant led to psychotic type state in user in Southern Trust area
10 th December	Increased levels of MDMA in ecstasy pills in a number of European countries
24 th December	Deaths in Sweden associated with Bonzai Citrus / Bonzai Winter Boost
5 th January	Superman pills containing lethal amounts of PMMA found in Netherlands, and linked to deaths in England
30 th January	Botulism linked to injecting drug users hospitalised in Scotland
9 th February	UPS ecstasy tablets seized in Europe found to contain very high levels of MDMA
10 th February	UPS tablets seized in a number of areas in Northern Ireland
18 th February	"Trippy weed" linked to hospitalisation of 2 girls in Southern Trust area
25 th February	2 people believed to have been hospitalised after smoking unknown substance (believed by users to be sky blue / sky high)
27 th February	Unknown oily liquid (possible GHB; GBL) led to unconsciousness in user
3 rd March	Heineken tablets in England believed to contain very high levels of MDMA
9 th March	Unknown bright yellow powder (believed by user to be mephedrone) causing strong adverse effects and leading to addiction
12 th March	Report of "blue sky" in Belfast area being very potent, and user having seizure
20 th March	Report of young person being admitted to Emergency Department with potentially life threatening symptoms after using "blue sky".

Information circulated by DAMIS during October 2014 – March 2015

In the majority of the above cases, more information was sought either from the original informant, or from other experts in the field (Addictions staff, Homeless staff and drug users) and DAMIS partners continue to monitor the situation. In some cases, information was sent to specific groups, for example GP Practices, or Needle Exchange Providers.

In some cases, information was considered to be of sufficient urgency for it to be circulated to the full DAMIS database. Information circulated to the full DAMIS database is shown in the table below:

Emails circulated by DAMIS to the full DAMIS database <i>October 2014 – March 2015</i>		
Date	Topic	Nature of email
13 th November	Sky blue / sky high alert and synthetic cannabinoids factsheet	Warning over synthetic cannabinoids
28 th November	White heroin Amsterdam	Warning over white heroin being sold as cocaine
2 December	Ecstasy	Ecstasy linked to severe comedowns and suicidal thoughts
2 January	Botulism	Cases of Botulism in injecting drug users in Glasgow
6 January	Superman logo tablets	May contain lethal doses of PMMA
10 th February	UPS logo tablets	May contain very high doses of MDMA
13 th March	Sky blue / sky high alert and synthetic cannabinoids factsheet	Warning over synthetic cannabinoids

Additional awareness raising

In addition to the DAMIS alert on botulism on 2nd January, the Chief Medical Officer issued a CMO Alert to key medical staff <http://www.dhsspsni.gov.uk/hss-md-1-2015.pdf>

Reports of UPS logo ecstasy pills containing very high levels of MDMA, along with a number of PSNI seizures of such tablets, were considered of sufficient urgency for the Public Health Agency to issue a press release on this topic on 12th February.

The PHA is in ongoing discussions with partners around the potential need for further awareness raising around synthetic cannabinoids.

Information Sheet

Overdoses & Emergencies



Date: 14/04/2014
Version: 1.0

Overdoses of depressant drugs often involve breathing difficulties, while overdoses of stimulant drugs can involve heart attacks or fits. Because of this, you may need to do different things to help someone. What you should do depends on their appearance and behaviour.



Vomiting/feeling unwell: vomiting is usually nature's way of telling you've had too much. If somebody is unwell, don't give them anything to eat and only let them drink water (never force them to drink anything). If after vomiting they want to sleep, let them but keep your eye on them. Make sure they are lying on their side (see the **recovery position** on next page.)



If they are overheating: cool them down by removing outer clothing; fan them; use a wet cloth on their skin*; take them outside or somewhere cool. If they are conscious allow them to sip water or a non alcoholic drink. **Call an ambulance.**

**Do not use very cold water, this can repel the superficial blood vessels deeper into the body and prevent heat loss. Even lukewarm water is fine as it mimics the temperature of sweat, the body's natural way to reduce temperature.*



If they have chest pains: sit them down in a calm environment and reassure them. **Call an ambulance.**



Bad trip/freak out/paranoia: if somebody is having a frightening or disturbing drug experience or have become very paranoid, take them somewhere that is quiet where they feel safe (ideally a low stimulus environment and not a dance floor in a nightclub). Try to calm and reassure them (*"it will pass - the drugs will wear off"*). This can take hours, so be prepared to be patient. If they become panicky and you notice them breathing very fast, get them to control their breathing by slowing it down or breathing into a paper bag. If any of these disturbing experiences carry on after the drug has worn off, they need to speak to a doctor or drug service.



Serotonin syndrome: Serotonin syndrome can kill if it is not dealt with quickly by **calling for an ambulance**. Serotonin syndrome is a result of your body releasing too much of the neurotransmitter serotonin. It can be triggered by a number of different drugs. The most severe cases involve interactions of drugs that release serotonin, such as MDMA (ecstasy) and a range of other drugs known as 'serotonin re-uptake inhibitors'.

The main symptoms of serotonin syndrome are: rigid, jerky, twitchy unusual movements, often involving the legs shaking; fully dilated pupils; overheating; shivering; racing heart; the person appearing agitated and confused. If in doubt, ring for an ambulance.

It is important if they have rigid, jerky movements, not to hold people down because of the risk of muscle tissue breaking down (*rhabdomyolysis*). As with people who have been using *volatile substances* (solvents) it can also be risky to startle or frighten people as this can lead to heart failure.



If they can't be woken: (by shaking their shoulders and calling their name), or you notice a blueness of the skin, including lips or fingernails (or greyish for darker complexions) or they have trouble breathing, **call an ambulance.**

Check breathing: try to assess the airway and then breathing. If there is no breathing or it is abnormal (e.g. death rattle, agonal breath) then **CPR** should be attempted.

Check there is nothing stuck in their throat (vomit etc), if there is remove it. For vomit turn the head to the side and let gravity do its job. If that doesn't work turn their far shoulder towards you so that their mouth points towards the ground for 5 secs. If neither work don't waste time, start CPR or they will die quickly.

CPR: this can be **chest compressions** alone. If you know how and feel able to, give 30 chest compressions followed by 2 **rescue breaths**. These compressions and rescue breaths are called 1 cycle of CPR and should be repeated.



If they are having a 'fit': make sure the area is safe and there is nothing they could hurt themselves on. **Call an ambulance.** Be sure to inform the paramedics if the fit stops and starts, if it doesn't stop within a couple of minutes or if the person turns blue.

If they are unconscious, but still breathing normally (at least 1 normal breath in a 10 second period) **put them in the recovery position and call an ambulance.**

see next page

"Look after people who have overdosed in the same way you would want them to look after you"

Information Sheet

Overdoses & Emergencies



Date: 14/04/2014
Version: 1.0

If somebody is unconscious and then vomits while lying on their back, they can swallow their vomit and literally drown in it. That is why you should put an unconscious person in the recovery position and **call for an ambulance**.

The Recovery Position



Put the hand closest to you by the head (as if they were waving)



Put the arm furthest away from you across the chest, so that the back of the hand rests against the cheek



Hold the hand in place and lift up the knee furthest away from you, making sure the foot is planted firmly on the ground



Turn them on their side by pushing down on their knee

Antidotes

Doctors and paramedics can administer an antidote to some types of overdoses caused by depressants. If it is an opiate (eg. heroin) overdose and there is **naloxone*** available you should administer it as directed by its Patient Information Leaflet within the naloxone pack. It is perfectly legal for you to do so in an emergency. t

*In some areas naloxone is given out as *Prenoxad*, a licensed product but still containing naloxone HCL (at 1mg/ml).

Calling an ambulance

Never hesitate to call an ambulance. In most areas, the police are only called to overdoses if there is a death or an under 16 involved, or if there has been a previous incident of violence at the address given. In some areas the police may also attend if the caller states that the casualty is not breathing normally or not breathing at all. In this circumstance their priority is the preservation of life rather than law enforcement.

"Look after people who have overdosed in the same way you would want them to look after you"